

# Biodiversity: Understanding Complex Ecological Systems as Bifurcations of the Species World Line

Alfred Russel Wallace

Ian Malcolm

Ternate Biodiversity Institute  
Ternate, Indonesia

University of Western Costa Rica  
and the Santa Fe Institute  
malcolm@santafe.edu

## Abstract

By utilising an observational approach we are able to contextualise the current debate on the origins of biodiversity, particularly the present amorphous definitions of species and their stability, and place these concepts in their proper emergent framework. Returning to first principles and characterising life as a “struggle for existence” we are able to show how speciation events should be considered as the bifurcation of a representational dynamical system i.e. an environmentally mediated feedback effect with long term consequences for the diversity of life on earth. Finally we consider the lessons to be learnt through studies of captive breeding programs and discuss the Lamarckian implications of recent advances in genetic knowledge *vis a vis* this paper.

To appear in *Posthumous Preprints in Emergence* 1:1 (1998)

## Contents

1	Introduction	1
2	The Struggle for Existence	2
3	The Law of Population of Species	2
4	The Abundance or Rarity of a Species dependent upon its more or less perfect Adaptation to the Conditions of Existence	4
5	Useful Variations will tend to Increase; useless or hurtful Variations to Diminish	5
6	Superior Varieties will ultimately Extirpate the original Species	6
7	The Partial Reversion of Domesticated Varieties explained	7
8	Lamarck's Hypothesis very different from that now advanced	8
9	Conclusion	9

## 1 Introduction

“No one ought to feel surprise at much remaining as yet unexplained in regard to the origin of species and varieties, if he makes due allowance for our profound ignorance in regard to the mutual relations of all the beings which live around us.”

from *On the Origin of Species*

— Charles Darwin, 1859

ONE of the strongest arguments which have been adduced to prove the original and permanent distinctness of species is, that varieties produced in a state of domesticity are more or less unstable, and often have a tendency, if left to themselves, to return to the normal form of the parent species; and this instability is considered to be a distinctive peculiarity of all varieties, even of those occurring among wild animals in a state of nature, and to constitute a provision for preserving unchanged the originally created distinct species.

In the absence of scarcity of facts and observations as to varieties occurring among wild animals, this argument has had great weight with naturalists, and has led to a very general and somewhat prejudiced belief in the stability of species. Equally general, however, is the belief in what are called “permanent or true varieties,” — races of animals which continually propagate their like, but which differ so slightly (although constantly) from some other race, that the one is considered to be a variety of the other. Which is the variety and which the original species, there is generally no means of determining, except in those rare cases in which the one race has been known to produce an offspring unlike

itself and resembling the other. This, however, would seem quite incompatible with the “permanent invariability of species,” but the difficulty is overcome by assuming that such varieties have strict limits, and can never again vary further from the original type, although they may return to it, which, from the analogy of the domesticated animals, is considered to be highly probable, if not certainly proved.

It will be observed that this argument rests entirely on the assumption, that varieties occurring in a state of nature are in all respects analogous to or even identical with those of domestic animals, and are governed by the same laws as regards their permanence or further variation. But it is the object of the present paper to show that this assumption is altogether false, that there is a general principle in nature which will cause many varieties to survive the parent species, and to give rise to successive variations departing further and further from the original type, and which also produces, in domesticated animals, the tendency of varieties to return to the parent form.

## **2 The Struggle for Existence**

The life of wild animals is a struggle for existence. The full exertion of all their faculties and all their energies is required to preserve their own existence and provide for that of their infant offspring. The possibility of procuring food during the least favourable seasons, and of escaping the attacks of their most dangerous enemies, are the primary conditions which determine the existence both of individuals and of entire species. These conditions will also determine the population of a species; and by a careful consideration of all the circumstances we may be enabled to comprehend, and in some degree to explain, what at first sight appears so inexplicable — the excessive abundance of some species, while others closely allied to them are very rare.

## **3 The Law of Population of Species**

The general proportion that must obtain between certain groups of animals is readily seen. Large animals cannot be so abundant as small ones; the carnivora must be less numerous than the herbivora; eagles and lions can never be so plentiful as pigeons and antelopes; the wild asses of the Tartarian deserts cannot equal in numbers the horses of the more luxuriant prairies and pampas of America. The greater or less fecundity of an animal is often considered to be one of the chief causes of its abundance or scarcity; but a consideration of the facts will show us that it really has little or nothing to do with the matter. Even the least prolific of animals would increase rapidly if unchecked, whereas it is evident that the animal population of the globe must be stationary, or perhaps, through the influence of man, decreasing. Fluctuations there may be; but permanent increase, except in restricted localities, is almost impossible.

For example, our own observation must convince us that birds do not go

on increasing every year in a geometrical ratio, as they would do, were there not some powerful check to their natural increase. Very few birds produce less than two young ones each year, while many have six, eight, or ten; four will certainly be below the average; and if we suppose that each pair produce young only four times in their life, that will also be below the average, supposing them not to die either by violence or want of food. Yet at this rate how tremendous would be the increase in a few years from a single pair! A simple calculation will show that in fifteen years each pair of birds would have increased to nearly ten millions! whereas we have no reason to believe that the number of the birds of any country increases at all in fifteen or in one hundred and fifty years. With such powers of increase the population must have reached its limits, and have become stationary, in a very few years after the origin of each species.

It is evident, therefore, that each year an immense number of birds must perish — as many in fact as are born; and as on the lowest calculation the progeny are each year twice as numerous as their parents, it follows that, whatever be the average number of individuals existing in any given country, twice that number must perish annually, — a striking result, but one which seems at least highly probable, and is perhaps under rather than over the truth. It would therefore appear that, as far as the continuance of the species and the keeping up the average number of individuals are concerned, large broods are superfluous. On the average all above one become food for hawks and kites, wild cats and weasels, or perish of cold and hunger as winter comes on.

This is strikingly proved by the case of particular species; for we find that their abundance in individuals bears no relation whatever to their fertility in producing offspring. Perhaps the most remarkable instance of an immense bird population is that of the passenger pigeon of the United States, which lays only one, or at most two eggs, and is said to rear generally but one young one. Why is this bird so extraordinarily abundant, while others producing two or three times as many young are much less plentiful? The explanation is not difficult. The food most congenial to this species, and on which it thrives best, is abundantly distributed over a very extensive region, offering such difference of soil and climate, that in one part or another of the area the supply never fails. The bird is capable of a very rapid and long-continued flight, so that it can pass without fatigue over the whole of the district it inhabits, and as soon as the supply of food begins to fail in one place is able to discover a fresh feeding-ground.

This example strikingly shows us that the procuring a constant supply of wholesome food is almost the sole condition requisite for ensuring the rapid increase of a given species, since neither the limited fecundity, nor the unrestrained attacks of birds of prey and of man are here sufficient to check it. In no other birds are these peculiar circumstances so strikingly combined. Either their food is more liable to failure, or they have not sufficient power of wing to search for it over an extensive area, or during some season of the year it becomes very scarce, and less wholesome substitutes have to be found; and thus, though more fertile in offspring, they can never increase beyond the supply of food in the least favourable seasons.

Many birds can only exist by migrating, when their food becomes scarce,

to regions possessing a milder, or at least a different climate, though, as these migrating birds are seldom excessively abundant, it is evident that the countries they visit are still deficient in a constant and abundant supply of wholesome food. Those whose organization does not permit them to migrate when their food becomes periodically scarce, can never attain a large population. This is probably the reason why woodpeckers are scarce with us, while in the tropics they are among the most abundant of solitary birds. Thus the house sparrow is more abundant than the redbreast, because its food is more constant and plentiful, — seeds of grasses being preserved during the winter, and our farm-yards and stubble-fields furnishing an almost inexhaustible supply. Why, as a general rule, are aquatic, and especially sea birds, very numerous in individuals? Not because they are more prolific than others, generally the contrary; but because their food never fails, the sea-shores and river-banks daily swarming with a fresh supply of small mollusca and crustacea.

Exactly the same laws will apply to mammals. Wild cats are prolific and have few enemies; why then are they never as abundant as rabbits? The only intelligible answer is, that their supply of food is more precarious. It appears evident, therefore, that so long as a country remains physically unchanged, the numbers of its animal population cannot materially increase. If one species does so, some others requiring the same kind of food much diminish in proportion. The numbers that die annually must be immense; and as the individual existence of each animal depends upon itself, those that die must be the weakest — the very young, the aged, and the diseased, — while those that prolong their existence can only be the most perfect in health and vigour — those who are best able to obtain food regularly, and avoid their numerous enemies. It is, as we commenced by remarking, “a struggle for existence,” in which the weakest and least perfectly organized must always succumb.

#### **4 The Abundance or Rarity of a Species dependent upon its more or less perfect Adaptation to the Conditions of Existence**

It seems evident that what takes place among the individuals of a species must also occur among the several allied species of a group, i.e. that those which are best adapted to obtain a regular supply of food, and to defend themselves against the attacks of their enemies and the vicissitudes of the seasons, must necessarily obtain and preserve a superiority in population; while those species which from some defect of power or organization are the least capable of counteracting the vicissitudes of food, supply, &c., must diminish in numbers, and, in extreme cases, become altogether extinct.

Between these extremes the species will present various degrees of capacity for ensuring the means of preserving life; and it is thus we account for the abundance or rarity of species. Our ignorance will generally prevent us from accurately tracing the effects to their causes; but could we become perfectly

acquainted with the organization and habits of the various species of animals, and could we measure the capacity of each for performing the different acts necessary to its safety and existence under all the varying circumstances by which it is surrounded, we might be able even to calculate the proportionate abundance of individuals which is the necessary result.

If now we have succeeded in establishing these two points — 1st, that the animal population of a country is generally stationary, being kept down by a periodical deficiency of food, and other checks; and, 2nd, that the comparative abundance or scarcity of the individuals of the several species is entirely due to their organization and resulting habits, which, rendering it more difficult to procure a regular supply of food and to provide for their personal safety in some cases than in others, can only be balanced by a difference in the population which have to exist in a given area — we shall be in a condition to proceed to the consideration of varieties, to which the preceding remarks have a direct and very important application.

## **5 Useful Variations will tend to Increase; useless or hurtful Variations to Diminish**

Most or perhaps all the variations from the typical form of a species must have some definite effect, however slight, on the habits or capacities of the individuals. Even a change of colour might, by rendering them more or less distinguishable, affect their safety; a greater or less development of hair might modify their habits. More important changes, such as an increase in the power or dimensions of the limbs or any of the external organs, would more or less affect their mode of procuring food or the range of country which they inhabit. It is also evident that most changes would affect, either favourably or adversely, the powers of prolonging existence. An antelope with shorter or weaker legs must necessarily suffer more from the attacks of the feline carnivora; the passenger pigeon with less powerful wings would sooner or later be affected in its powers of procuring a regular supply of food; and in both cases the result must necessarily be a diminution of the population of the modified species. If, on the other hand, any species should produce a variety having slightly increased powers of preserving existence, that variety must inevitably in time acquire a superiority in numbers.

These results must follow as surely as old age, intemperance, or scarcity of food produce an increased mortality. In both cases there may be many individual exceptions; but on the average the rule will invariably be found to hold good. All varieties will therefore fall into two classes — those which under the same conditions would never reach the population of the parent species, and those which would in time obtain and keep a numerical superiority. Now, let some alteration of physical conditions occur in the district — a long period of drought, a destruction of vegetation by locusts, the irruption of some new carnivorous animal seeking “pastures new” — any change in fact tending to render existence more difficult to the species in question, and tasking its utmost

powers to avoid complete extermination; it is evident that, of all the individuals composing the species, those forming the least numerous and most feebly organized variety would suffer first, and, were the pressure severe, must soon become extinct. The same causes continuing in action, the parent species would next suffer, would gradually diminish in numbers, and with a recurrence of similar unfavourable conditions might also become extinct. The superior variety would then alone remain, and on a return to favourable circumstances would rapidly increase in numbers and occupy the place of the extinct species and variety.

## **6 Superior Varieties will ultimately Extirpate the original Species**

The variety would now have replaced the species, of which it would be a more perfectly developed and more highly organized form. It would be in all respects better adapted to secure its safety, and to prolong its individual existence and that of the race. Such a variety could not return to the original form; for that form is an inferior one, and could never compete with it for existence. Granted, therefore, a “tendency” to reproduce the original type of the species, still the variety must ever remain preponderant in numbers, and under adverse physical conditions again alone survive. But this new, improved, and populous race might itself, in course of time, give rise to new varieties, exhibiting several diverging modifications of form, any of which, tending to increase the facilities for preserving existence, must by the same general law, in their turn become predominant.

Here, then, we have progression and continued divergence deduced from the general laws which regulate the existence of animals in a state of nature, and from the undisputed fact that varieties do frequently occur. It is not, however, contended that this result would be invariable; a change of physical conditions in the district might at times materially modify it, rendering the race which had been the most capable of supporting existence under the former conditions now the least so, and even causing the extinction of the newer and, for a time, superior race, while the old or parent species and its first inferior varieties continued to flourish. Variations in unimportant parts might also occur, having no perceptible effect on the life-preserving powers; and the varieties so furnished might run a course parallel with the parent species, either giving rise to further variations or returning to the former type.

All we argue for is, that certain varieties have a tendency to maintain their existence longer than the original species, and this tendency must make itself felt; for though the doctrine of chances or averages can never be trusted to on a limited scale, yet, if applied to high numbers, the results come nearer to what theory demands, and, as we approach to an infinity of examples, become strictly accurate. Now the scale on which nature works is so vast — the numbers of individuals and periods of time with which she deals approach so near to infinity, that any cause, however slight, and however liable to be veiled and counteracted

by accidental circumstances, must in the end produce its full legitimate results.

## 7 The Partial Reversion of Domesticated Varieties explained

Let us now turn to domesticated animals, and inquire how varieties produced among them are affected by the principles here enunciated. The essential difference in the condition of wild and domestic animals is this, — that among the former, their well-being and very existence depend upon the full exercise and healthy condition of all their senses and physical powers, whereas, among the latter, these are only partially exercised, and in some cases are absolutely unused. A wild animal has to search, and often to labour, for every mouthful of food — to exercise sight, hearing, and smell in seeking it, and in avoiding dangers, in procuring shelter from the inclemency of the seasons, and in providing for the subsistence and safety of its offspring. There is no muscle of its body that is not called into daily and hourly activity; there is no sense or faculty that is not strengthened by continual exercise.

The domestic animal, on the other hand, has food provided for it, is sheltered, and often confined, to guard it against the vicissitudes of the seasons, is carefully secured from the attacks of its natural enemies, and seldom even rears its young without human assistance. Half of its senses and faculties are quite useless; and the other half are but occasionally called into feeble exercise, while even its muscular system is only irregularly called into action.

Now when a variety of such an animal occurs, having increased power or capacity in any organ or sense, such increase is totally useless, is never called into action, and may even exist without the animal ever becoming aware of it. In the wild animal, on the contrary, all its faculties and powers being brought into full action for the necessities of existence, any increase becomes immediately available, is strengthened by exercise, and must even slightly modify the food, the habits, and the whole economy of the race. It creates as it were a new animal, one of superior powers, and which will necessarily increase in numbers and outlive those inferior to it.

Again, in the domesticated animal all variations have an equal chance of continuance; and those which would decidedly render a wild animal unable to compete with its fellows and continue its existence are no disadvantage whatever in a state of domesticity. Our quickly fattening pigs, short-legged sheep, pouter pigeons, and poodle dogs could never have come into existence in a state of nature, because the very first step towards such inferior forms would have led to the rapid extinction of the race; still less could they now exist in competition with their wild allies. The great speed but slight endurance of the race horse, the unwieldy strength of the ploughman's team, would both be useless in a state of nature. If turned wild on the pampas, such animals would probably soon become extinct, or under favorable circumstances might each lose those extreme qualities which would never be called into action, and in a few generations would

revert to a common type, which must be that in which the various powers and faculties are so proportioned to each other as to be best adapted to procure food and secure safety, — that in which by the full exercise of every part of his organization the animal can alone continue to live. Domestic varieties, when turned wild, must return to something near the type of the original wild stock, or become altogether extinct<sup>1</sup>.

## 8 Lamarck's Hypothesis very different from that now advanced

We see, then, that no inferences as to varieties in a state of nature can be deduced from the observation of those occurring among domestic animals. The two are so much opposed to each other in every circumstance of their existence, that what applies to the one is almost sure not to apply to the other. Domestic animals are abnormal, irregular, artificial; they are subject to varieties which never occur and never can occur in a state of nature; their very existence depends altogether on human care: so far are many of them removed from that just proportion of faculties, that true balance of organization, by means of which alone an animal left to its own resources can preserve its existence and continue its race.

The hypothesis of Lamarck — that progressive changes in species have been produced by the attempts of animals to increase the development of their own organs, and thus modify their structure and habits — has been repeatedly and easily refuted by all writers on the subject of varieties and species, and it seems to have been considered that when this was done the whole question has been finally settled; but the view here developed renders such an hypothesis quite unnecessary, by showing that similar results must be produced by the action of principles constantly at work in nature.

The powerful retractile talons of the falcon — and the cat-tribes have not been produced or increased by the volition of those animals; but among the different varieties which occurred in the earlier and less highly organized forms of these groups, those always survived longest which had the greatest facilities for seizing their prey. Neither did the giraffe acquire its long neck by desiring to reach the foliage of the more lofty shrubs, and constantly stretching its neck for the purpose, but because any varieties which occurred among its antitypes with a longer neck than usual at once secured a fresh range of pasture over the same ground as their shorter-necked companions, and on the first scarcity of food were thereby enabled to outlive them.

Even the peculiar colours of many animals, especially insects, so closely resembling the soil or the leaves or the trunks on which they habitually reside, are explained on the same principle; for though in the course of ages varieties of many tints may have occurred, yet those races having colours best adapted

---

<sup>1</sup>That is, they will vary, and the variations which tend to adapt them to the wild state, and therefore approximate them to wild animals, will be preserved. Those individuals which do not vary sufficiently will perish.

to concealment from their enemies would inevitably survive the longest. We have also here an acting cause to account for that balance so often observed in nature, — a deficiency in one set of organs always being compensated by an increased development of some others — powerful wings accompanying weak feet, or great velocity making up for the absence of defensive weapons; for it has been shown that all varieties in which an unbalanced deficiency occurred could not long continue their existence.

The action of this principle is exactly like that of the centrifugal governor of the steam engine, which checks and corrects any irregularities almost before they become evident; and in like manner no unbalanced deficiency in the animal kingdom can ever reach any conspicuous magnitude, because it would make itself felt at the very first step, by rendering existence difficult and extinction almost sure to follow. An origin such as is here advocated will also agree with the peculiar character of the modifications of form and structure which obtain in organized beings — the many lines of divergence from a central type, the increasing efficiency and power of a particular organ through a succession of allied species, and the remarkable persistence of unimportant parts such as colour, texture of plumage and hair, form of horns or crests, through a series of species differing considerably in more essential characters. It also furnishes us with a reason for that “more specialized structure” which Professor Owen states to be a characteristic of recent compared with extinct forms, and which would evidently be the result of the progressive modification of any organ applied to a special purpose in the animal economy.

## 9 Conclusion

We believe we have now shown that there is a tendency in nature to the continued progression of certain classes of varieties further and further from the original type — a progression to which there appears no reason to assign any definite limits — and that the same principle which produces this result in a state of nature will also explain why domestic varieties have a tendency to revert to the original type. This progression, by minute steps, in various directions, but always checked and balanced by the necessary conditions, subject to which alone existence can be preserved, may, it is believed, be followed out so as to agree with all the phenomena presented by organized beings, their extinction and succession in past ages, and all the extraordinary modifications of form, instinct, and habits which they exhibit.